

Product no **AS12 1871**

ABI2 | Absciscic acid insensitive 2

Product information

Background	Protein Absciscic acid-insensitive 2 (ABI2) is a represor of the ABA signaling pathway, regulating various ABA responses like: stomatal closure, osmotic water permability of the plasma membrane, high light stress, response to glucose, seed germination and inhibition of vegetative growth. Involved in acquired thermotolerance. Alternative names: ABA INSENSITIVE 2, ABI2, ATABI2, Protein phosphatase 2C 77, AtPP2C77, At5g57050, MHM17_19, MHM17.19, PP2C ABI2.
Immunogen	<u>KLH</u> -conjugated synthetic peptide derived from N-terminus of <i>Arabidopsis thaliana</i> ABI2 sequence, UniProt: O04719 , TAIR: AT5G57050 chosen peptide is not conserved in ABI1
Host	Rabbit
Clonality	Polyclonal
Purity	Affinity purified serum
Format	Lyophilized in PBS pH 7.4
Quantity	50 µg
Reconstitution	For reconstitution add 50 µl of sterile water.
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tubes.
Tested applications	Western blot (WB)
Related products	AS12 1861 Anti-ABI1 absciscic acid insensitive 1, rabbit antibodies AS12 1862 Anti-ABI3 absciscic acid insensitive 3, rabbit antibodies AS12 1863 Anti-ABI5 absciscic acid insensitive 5, rabbit antibodies AS13 2634 Anti-PYR1 Absciscic acid receptor RCAR11, rabbit antibodies Plant protein extraction buffer Secondary antibodies

Application information

Expected apparent MW	46 kDa
Confirmed reactivity	<i>Arabidopsis thaliana</i>
Not reactive in	No confirmed exceptions from predicted reactivity are currently known.
Additional information	<p>ABI2 antibodies recognize recombiant StrepTag-ABI2, ABI2-GST, His-ABI2. ABI2 protein is easily degraded therefore extraction buffer needs to contain protease inhibitors, example of such inhibitor cocktail can be found here.</p> <p>To detect endogenous ABI2 plant material needs to be subjected to stress before harvesting. For high resolution images, please visit the specific product page at www.agrisera.com</p>
Selected references	Mitula et al. (2015). Arabidopsis ABA-Activated Kinase MAPKKK18 is Regulated by Protein Phosphatase 2C ABI1 and the Ubiquitin-Proteasome Pathway. Plant Cell Physiol. 2015 Dec;56(12):2351-67. doi: 10.1093/pcp/pcv146. Epub 2015 Oct 6.

For high resolution images, please visit the specific product page at www.agrisera.com


Agrisera

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

application example



GST-ABI1 and GST-ABI2 were purified on glutathione sepharose and separated on 10% SDS-PAGE and blotted 1h to PVDF (semi-dry). Blots were blocked with 3% semi-skimmed milk for 30 min. at room temperature (RT) with agitation. Blots were incubated with the anti-ABI2 primary antibody diluted to 1: 1000 for 30 minutes at RT with agitation. The antibody solution was decanted and the blot was rinsed briefly twice, then washed 3 times for 5 min in TBS-T at RT with agitation. Blot was incubated in secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated, from Agrisera, [AS09 602](#)) diluted to 1:50 000 for 30 min. at RT with agitation. The blot was washed as above and developed for 1 min with ECL according to the manufacturer's instructions. Exposure time was 5 min. Multiple bands are a result of degradation of GST-ABI2 protein.

Courtesy of Małgorzata Tajdel, from Dr. Agnieszka Ludwików laboratory, Adam Mickiewicz University, Poland